

UAHEP986
IFP-764-UNC
November 1998
hep-ph/9811326

Neutrino Masses in R Parity Violating Supersymmetry

L. Clavelli* and P.H. Frampton †

**Department of Physics and Astronomy, University of Alabama,
Tuscaloosa AL 35487*

*† Department of Physics and Astronomy, University of North Carolina,
Chapel Hill NC 27599-3255*

Abstract

We discuss a model for neutrino masses and mixings based on R parity violating Yukawa couplings. The model requires no right-handed neutrinos. It accommodates the SuperKamiokande data on atmospheric neutrinos and successfully relates this data to the mass difference for solar neutrinos in the small-angle MSW solution. We obtain an unexpected testable pattern for the three neutrino flavors.

PACS numbers: 12.60.Jv, 12.38.Bx, 14.60.Pq, 14.80.Ly

Typeset using REVTeX

*lclavell@bama.ua.edu

†frampton@physics.unc.edu

Recently the SuperKamiokande collaboration [1] presented striking evidence for neutrino masses and mixings. The data extend by more than six orders of magnitude the previously known hierarchy among fermion masses. The result could suggest the existence of right handed neutrinos as required, for example, in models of leptons as a fourth color [2] or in left-right symmetric models [3]. The extreme smallness of the neutrino masses might afford a window on the grand unification scale [4] through the see-saw mechanism [5,6]. The importance of these questions makes it imperative to give wide latitude to the discussion of alternative possibilities for understanding neutrino masses. The literature is, as expected, mushrooming [7–20].

The Class of Models. In this paper we consider R parity violating (RPV) supersymmetry-based models leading to calculable neutrino masses. We arrive at a specific pattern of masses and mixings quite different from the see-saw mechanism. Similar models have already been the subject of study by other authors [21–25].

For definiteness we work in the context of gravity mediated supersymmetry (SUSY) breaking scheme (MSSM) with additional RPV Yukawa couplings. Gauge mediated SUSY breaking is also compatible with the mechanism we wish to discuss. After this introduction we discuss a simple ansatz that leads to the observed neutrino squared mass differences and the apparent large mixing seen in the atmospheric experiments, and discuss a slight generalization of the simplest model which retains the same general features. Then, we describe a Monte-Carlo search in the parameter space of Yukawa couplings which favors this type of model and hence increases our confidence that our choices are not arbitrary or *ad hoc*. There follow the general constraints on the RPV couplings arising from the neutrino observations, together with a summary and some pertinent general observations.

The One-Loop Graph. For calculability of neutrino masses we assume the absence of right handed neutrinos and tree level Majorana masses. The latter would be forbidden, for example, in a model where there are no dimension three operators in the SUSY breaking Lagrangian. Such a model would also forbid gluino and photino masses at tree level and leaves open the interesting possibility that the neutrinos could be mixing with the photino. The minimum expectation for the photino mass, however, would be in the MeV region far above the current range of apparent neutrino masses so we do not consider this possible component in the mixing matrix at the present time. Whether or not the gluino and photino are light has no direct bearing on the current model. We, therefore, do not discuss proposed counter-indications to the light gluino scenario from LEP four jet analyses [26] and counter-arguments thereto [27]. Similarly the LEP II indications from Chargino and Higgs searches that the Higgs sector in the MSSM might have to be enlarged by a singlet Higgs boson in the $m_{1/2} = A = 0$ model is not directly relevant to our present considerations.

We consider a Lepton number and R parity violating term in the superpotential of the form

$$\mathcal{W} = \lambda_{ijk} e L^i L^j \bar{E}^k \quad (1)$$




FIG. 1. R-parity violating graph leading to neutrino mixing

where L represents the left handed lepton doublet superfield in the i 'th family and E represents the singlet lepton superfield of the k 'th family. The renormalization group behavior of this RPV coupling is discussed in [28]. Because of the superfield antisymmetry, the λ 's are antisymmetric in the first two indices leaving nine independent components. We have normalized them relative to the electric charge e . The corresponding piece of the interaction Lagrangian that will contribute to neutrino masses at one loop order is

$$\mathcal{L} = \lambda_{ijk} e (\tilde{e}_L^j \tilde{e}_R^k \nu_L^i + \tilde{e}_L^{j*} (\bar{\nu}_L^i) e_L^k - i \leftrightarrow j) \quad (2)$$

The loop graph of figure 1 yields the neutrino mass matrix

$$\mathcal{M}_{ij} = \sum_{k,l} \lambda_{ikl} \lambda_{jkl}^* \frac{\alpha}{3\pi} F(k, l) \quad (3)$$

The function $F(k, l)$ is the same as appears in the one loop contribution to gluino and photino masses [29]. It can be written in terms of the k 'th family charged lepton and slepton masses $m_k, \tilde{m}_{k,R}, \tilde{m}_{k,L}$ or in terms of the ratios $x_{k,L} = \frac{m_k^2}{\tilde{m}_{k,L}^2}, x_{k,R} = \frac{m_k^2}{\tilde{m}_{k,R}^2}$.

$$F(k, l) = m_l ((1 - x_{k,R})^{-1} \ln(x_{k,R}) - (1 - x_{k,L})^{-1} \ln(x_{k,L})) \quad (4)$$

We take the λ_{ijk} to be real for the purposes of this paper, ignoring possible CP violating phases. For the charged slepton masses we use the supergravity related expectations

$$\tilde{m}_{k,L}^2 = m_0^2 + m_k^2 + M_Z^2 \cos 2\beta [-1/2 + \sin^2 \theta] \quad (5)$$

$$\tilde{m}_{k,R}^2 = m_0^2 + m_k^2 - M_Z^2 \cos 2\beta \sin^2 \theta \quad (6)$$

Then, neglecting higher order terms, $F(k, l)$ takes the form

$$F(k, l) = \frac{M_Z^2}{M^2} \cos 2\beta (1/2 - \sin^2 \theta) m_l (1 + \frac{m_k^2}{M^2} \ln \frac{m_k^2}{M^2} + \mathcal{O}(1/M^4)) \quad (7)$$

where

$$M^2 = m_0^2 - \cos(2\beta)M_Z^2/4 \quad (8)$$

To a good approximation the neutrino masses are proportional to the the SUSY parameters in the combination $\lambda^2 \cos(2\beta)M_Z^2/M^2$. We adopt the nominal values $\tan(\beta) = 1.5, m_0 = 130\text{GeV}$ but other possible values can be accommodated by rescaling the λ 's by the appropriate factor.

The Simplest Model. The *simplest* model discussed in this paper is that where $\lambda_{121} = \lambda_{131} = \lambda_0$ and all other linearly independent λ 's vanish. With this assumption which will be further justified below one can fit the apparent neutrino squared mass differences suggested by the atmospheric and solar neutrino data and a large mixing in the $\nu_\mu - \nu_\tau$ channel with moderate values of λ_0 and no right handed neutrinos or GUT scale parameters. As discussed below, a discrete symmetry yields this pattern of the λ 's. As we shall see, whether or not the neutrino anomalies are due to these RPV couplings, the current data provide strong experimental constraints on the values of the independent λ 's.

We first note that the main features of the neutrino data can be fit in a simple one parameter model where this parameter is of moderate size. The model can be described by the as yet unmotivated assumption that all the λ 's of Eq.(1) vanish unless two of the three indices lie in the first generation and that these non-vanishing λ 's have equal magnitude. That is we assume

$$\lambda_{121} = \lambda_{131} = \lambda_0; \quad \lambda_{ijk} = 0 \quad (otherwise) \quad (9)$$

This pattern can be obtained by imposing a Z_2 symmetry on the superpotential. Under this Z_2 , the right-handed superfields and the first generation left-handed superfield changes sign. In order to provide for charged lepton masses there must then be at least two Higgs doublets, one of which couples to the first generation leptons and is even under the discrete symmetry while the other(s), coupling to the second and third generation leptons, are odd under the symmetry.

The non-zero elements of the neutrino mass matrix are now:

$$M_{11} = \frac{\alpha\lambda_0^2}{3\pi}(F(2,1) + F(3,1)) \quad (10)$$

and

$$M_{22} = M_{33} = M_{23} = M_{32} = \frac{\alpha\lambda_0^2}{3\pi}F(1,1) \quad (11)$$

In this zeroth order model, it is apparent that one cannot describe the mixing of the first generation (electron) neutrino. This mixing must be attributed to other interactions or to residual effects from other (small) λ' s. However, one can immediately see that this mass matrix incorporates a 45 deg mixing between the second and third generation neutrinos and we can show that the squared mass differences between the $\nu_\mu - \nu_\tau$ eigenstates and between the ν_e and the heavier of the other two neutrinos are consistent with observations for order unity values of λ_0 .

Diagonalizing the above mass matrix, the three eigenvalues are given by

$$m_1 = M_{11} \quad , \quad m_2 = 2M_{33} \quad , \quad m_3 = 0 \quad (12)$$

Substituting the expressions for $F(k, l)$ and neglecting non-leading powers of the lepton masses relative to slepton masses, the eigenstates have the squared mass differences

$$m_3^2 - m_2^2 = \left(\frac{\alpha \lambda_0^2}{3\pi} \left(\frac{m_Z^2}{M^2} \right) (1 - 4 \sin^2 \theta) \cos 2\beta \right)^2 m_e^2 \quad (13)$$

and

$$m_2^2 - m_1^2 = (m_3^2 - m_2^2) \left(\frac{m_\tau^2}{M^2} \ln \frac{m_\tau^2}{M^2} \right) \quad (14)$$

For example the two squared mass splittings are given consistent with experiment by reasonable SUSY parameters, $m_0 = 130$ GeV, $\cos(2\beta) = -.38$, if λ_0 is between .1 and 0.12. If m_0 ranges up to one TeV and $\tan\beta$ decreases slightly toward unity, the required λ_0 approaches unity. Independent of λ_0 the model predicts a ratio of the two squared mass differences in Eq.(14) consistent with the SuperKamiokande data and the MSW interpretation of the solar data for the universal scalar mass m_0 in the 100 GeV to 1 TeV range. The model disagrees with the vacuum mixing solution of the solar neutrino anomaly since this would require a much smaller value of the squared mass splitting ratio.

An Extension of the Simplest Model. As an extension of the model, we obtain the same mass matrix at leading order by allowing the more general set of relationships:

$$\lambda_{121} = \lambda_{131} \quad (15)$$

$$\lambda_{122} = \lambda_{132} \quad (16)$$

$$\lambda_{123} = \lambda_{133} \quad (17)$$

$$\lambda_{23k} = 0 \quad (18)$$

It is straightforwardly checked that this maintains the results of the previous subsection with the identification

$$\lambda_0 = \sum_k \lambda_{12k} m_k / m_e \quad (19)$$

General Search of All Possible λ_{ijk} Here we discuss a general search in the parameter space of the λ_{ijk} for values consistent with the atmospheric and solar neutrino experiments. Clearly, in order to fit the mass splitting observations and the apparent large mixing angle between the second and third eigenstates, the λ 's must lie in the neighborhood of the present simple model. However, in order to obtain non zero mixing of the electron neutrino from the Yukawa couplings, some small admixtures of the other λ 's are required. The allowed solutions are bounded by a nine dimensional rectangular parallelopiped whose side lengths we find by a Monte-Carlo method.

We impose the antisymmetry requirement in the first two indices on the λ_{ijk} and, running over all possible values of the nine independent components, we determine the neutrino mass matrix elements from Eq.(3).

Keeping the λ_{ijk} real, the mass matrix is diagonalized by an orthogonal matrix U satisfying

$$\mathcal{M}U = U\mathcal{M}_d \quad (20)$$

where $\mathcal{M}_d$ is the diagonal matrix of the three neutrino mass eigenstates. With no right handed neutrinos and with a large mixing in the $\nu_\mu - \nu_\tau$ sector, there is no possibility of finding a large mixing angle to ν_e . Thus we seek a solution corresponding to the small angle MSW solution. A set of nine λ_{ijk} is counted as a solution if it satisfies the following experimental constraints

- (1) $.0005eV^2 < |m_2^2 - m_3^2| < .006eV^2$
- (2) $.5365 < |U_{22}| < .843$ ($\sin(2\theta) > 0.83$)
- (3) $|U_{21}| < 0.1\sqrt{1 - U_{22}^2}$ (little mixing between ν_μ and ν_e)
- (4) $4 \cdot 10^{-6}eV^2 < |m_1^2 - m_3^2| < 1 \cdot 10^{-5}eV^2$ (MSW solution)
- (5) $0.9985 < |U_{11}| < 0.99985$ (small angle MSW case)

Without loss of generality we can choose one of the λ 's to be positive. We therefore choose $\lambda_{121} > 0$. Similarly, reversing the sign of all the λ 's with an odd number of index three's, has the effect of interchanging the eigenstates in the 2 – 3 sector. We therefore choose $\lambda_{131} > 0$ and label the heavier eigenstate in the strongly mixed 2 – 3 sector as ν_2 .

Constraints on the RPV Interactions. Here we compare the above constraints with RPV constraints found in earlier studies [25,30]. The solutions to the five constraints listed above require

$$0.0 < \lambda_{121} < 0.107 \quad (21)$$

$$-5.7 \cdot 10^{-3} < \lambda_{122} < 6.7 \cdot 10^{-3} \quad (22)$$

$$-1.9 \cdot 10^{-3} < \lambda_{123} < 1.6 \cdot 10^{-3} \quad (23)$$

$$-1.9 \cdot 10^{-2} < \lambda_{231} < 1.6 \cdot 10^{-2} \quad (24)$$

$$-8.8 \cdot 10^{-4} < \lambda_{232} < 1.2 \cdot 10^{-3} \quad (25)$$

$$-2.4 \cdot 10^{-4} < \lambda_{233} < 2.1 \cdot 10^{-4} \quad (26)$$

$$0.0 < \lambda_{131} < .106 \quad (27)$$

$$-8.0 \cdot 10^{-3} < \lambda_{312} < 6.9 \cdot 10^{-3} \quad (28)$$

$$-2.0 \cdot 10^{-3} < \lambda_{313} < 1.7 \cdot 10^{-3} \quad (29)$$

The upper limit on λ_{121} obtained here is less than the upper limit quoted earlier in the simple model due to the incorporation here of the electron neutrino mixing angle constraints.

If the neutrino mixing is primarily due to some mechanism other than that treated here, our results can be taken as upper limits on the λ 's with the results changing as discussed above if the other SUSY parameters are varied. The upper limits on the λ 's obtained here are comparable to the upper limits obtained from other constraints such as from charged current universality and from τ decay [30,31]. Thus the R-parity violating mechanism for neutrino mixing will give interesting predictions for other areas of phenomenology with only slight improvements in accuracy. For instance the upper limit of 29 on λ_{121} consistent with the neutrino data is close to the upper limit 0.17 (in our normalization) obtained in [30,31] from charged current universality.

Summary. Our main result is that a simple RPV pattern of couplings gives rise to the $\nu_\mu - \nu_\tau$ mixing with near maximal mixing consistent with the recent SuperKamiokande data. Scaling that mass difference to the experimentally- favored one of $\Delta m_{\mu\tau}^2 \sim 0.0025 eV^2$ and putting in the mixing angle for $\nu_e - \nu_\mu$ of $\sin^2 2\theta_{e\mu} \sim 0.01$ suggested by the small-angle MSW effect, one is led to a squared mass difference of $\Delta m_{e\mu}^2 \sim 10^{-5} eV^2$. A distinctive prediction of the current picture is that the mass of the electron neutrino is close to the heavier of the other two neutrino masses whereas most other models predict it closer to the lighter of the other two.

In fact the neutrino mass spectrum we predict is *upside-down* with respect to the see-saw expectation. Arbitrarily normalizing the mass eigenstate ν'_μ to $0.050000 eV$, our spectrum in the mass basis looks like: $M(\nu'_e) \simeq 0.050125 eV$; $M(\nu'_\mu) = 0.050000 eV$; and $M(\nu'_\tau) \simeq -0.000125 eV$. The mixing angles are approximately $\Theta_{\mu\tau} \simeq \pi/4$, $\Theta_{e\mu} \simeq 0.05$ and, although it is not fully constrained the third angle should be small $\Theta_{e\tau} \simeq \Theta_{\mu\tau} \simeq 0.05$. Of course, our subscripts for ν'_μ and ν'_τ in the mass basis are only suggestive, because both are approximately equal mixtures of ν_μ and ν_τ .

With this convention, the pattern which emerges for the squared neutrino mass differences is:

$$\Delta M_{\mu\tau}^2 \simeq \Delta M_{\tau e}^2 \gg \Delta M_{e\mu}^2 \quad (30)$$

while, at the same time, the neutrino mixing angles satisfy:

$$\Theta_{\mu\tau} \gg \Theta_{\tau e} \simeq \Theta_{e\mu}. \quad (31)$$

It would be interesting if this prediction for the neutrino masses and mixings is borne out by further experiments. It is possible that Eq.(30) and Eq.(31) are more general than this model.

Finally we re-emphasize that we are assuming only the minimal three left-handed neutrino fields ν_{iL} of the Standard Model. Also, we note that the electron neutrino mass $M(\nu_e)$, while two orders of magnitude beyond current bounds from the tritium β - decay end point, is only one order of magnitude beyond the present empirical limit arising from $(\beta\beta)_{0\nu}$, neutrinoless double β - decay.

This work was supported in part by the US Department of Energy under Grant no. DE-FG02-96ER-40967 at the University of Alabama and under Grant no. DE-FG05-85ER-40219 at the University of North Carolina.

REFERENCES

- [1] The Superkamiokande Collaboration, *hep-ex/9805006*. *hep-ex/9807003*.
Y. Fukuda et al. Phys. Rev. Lett. **81**, 1562 (1998)
- [2] J.C. Pati and A. Salam, Phys. Rev. Lett. **31**, 661 (1973)
- [3] R.N. Mohapatra and J.C. Pati, Phys. Rev. **D11**, 566 (1974)
- [4] J.K. Elwood, N. Irges, and P. Ramond, *hep-ph/9807228*.
- [5] M. Gell-Mann, P. Ramond, and R. Slansky, in Proceedings of the Workshop on Unified Theory and Baryon Number of the Universe, ed. T. Yanagida, KEK Japan, 1979
- [6] T. Yanagida, Prog. Theor. Phys. **B135**, 66 (1978).
- [7] M. Tanimoto, *hep-ph/9807283*.
- [8] G. Altarelli and F. Feruglio, Phys. Lett. **B439**, 112 (1998). *hep-ph/9807373*.
- [9] E. Ma, *hep-ph/9807386*.
- [10] H. Fritzsch and Z.-Z. Xing, *hep-ph/9807234*.
- [11] S. Lola and J.D. Vergados, Prog. Part. Nucl. Phys. **40**, 71 (1998). *hep-ph/9808269*.
- [12] R. Barbieri, L.J. Hall and A. Strumia, *hep-ph/9808333*.
- [13] R. Barbieri, L.J. Hall, D. Smith, A. Strumia and N. Weiner, *hep-ph/9807235*.
- [14] T. Teshima and T. Sakai, *hep-ph/9805386*.
- [15] K. Kang, S.K. Kang, C.S. Kim and S.M. Kim, *hep-ph/9808419*.
- [16] A. Joshipura, *hep-ph/9808261*.
- [17] J.C. Pati, *hep-ph/9807315*.
- [18] Y. Nomura and T. Yanagida, *hep-ph/9807325*.
- [19] P.H. Frampton, P.I. Krastev and J.T. Liu, Mod. Phys. Lett. **A9**, 761 (1994).
- [20] P. Langacker, Phys. Rev. **D58**, 093017 (1998). *hep-ph/9805281*.
- [21] F. Borzumati, Y. Grossman, E. Nardi and Y. Nir, Phys. Lett. **B384**, 123 (1996). *hep-ph/9606251*.
- [22] E.J. Chun, S.K. Kang, C.W. Kim, and U.W. Lee, *hep-ph/9807327*.
- [23] O.C.W. Kong, *hep-ph/9808304*.
- [24] B. Mukhopadhyaya, S. Roy and F. Vissani, *hep-ph/9808265*.
- [25] V. Barger, W.-Y. Keung and R.J.N. Phillips, Phys. Lett. **B364**, 27 (1995). *hep-ph/9507426*.
- [26] ALEPH Collaboration, Z. Phys. **C76**, 1 (1997)
- [27] G. Farrar, *hep-ph/9707467*.
- [28] V. Barger, M.S. Berger, R.J.N. Phillips, and T. Wohrmann, Phys. Rev. **D53**, 6407 (1996). *hep-ph/9511473*.
- [29] R. Barbieri, L. Girardello, and A. Masiero, Phys. Lett. **127B**, 429 (1983)
- [30] V. Barger, G.F. Giudice and T. Han, Phys. Rev. **D40**, 2987 (1989).
- [31] M. Carena, D. Choudhury, S. Lola, and C. Quigg, Phys. Rev. **D58**, 095003 (1998)